


On the Maximal Number of Real Embeddings of Spatial Minimally Rigid Graphs

Vangelis Bartzos, Ioannis Z. Emiris, Jan Legerský, Elias Tsigaridas
July 2018

The International Symposium on Symbolic and Algebraic Computation, New York





Rigidity in \mathbb{R}^3

An embedding $\rho: V \rightarrow \mathbb{R}^D$ of a graph $G = (V, E)$ is *compatible with edge lengths* $(d_{ij})_{ij \in E}$ if

$$\|\rho(i) - \rho(j)\| = d_{ij} \quad \text{for all } ij \in E.$$

Definition

A graph is *generically rigid* if the number of embeddings compatible with edge lengths induced by a generic embedding is finite modulo rotations and translations.

Rigidity in \mathbb{R}^3

An embedding $\rho: V \rightarrow \mathbb{R}^D$ of a graph $G = (V, E)$ is *compatible with edge lengths* $(d_{ij})_{ij \in E}$ if

$$\|\rho(i) - \rho(j)\| = d_{ij} \quad \text{for all } ij \in E.$$

Definition

A graph is *generically rigid* if the number of embeddings compatible with edge lengths induced by a generic embedding is finite modulo rotations and translations.

- If G is rigid and $G - \{e\}$ is not rigid $\forall e \in E$, then G is *minimally rigid*.
- $D = 2$: Laman graphs
(Capco, Gallet, Grasegger, Koutschan, Lubbes, Schicho, 2018)
- $D = 3$: Geiringer graphs

Algebraic Equations

Fix coordinates of a triangle to remove rigid motions.

$$(x_i - x_j)^2 + (y_i - y_j)^2 + (z_i - z_j)^2 = d_{ij}^2 \text{ for } ij \in E$$

- $\#$ complex solutions is an upper bound
- Loose mixed volume bound

Algebraic Equations

Fix coordinates of a triangle to remove rigid motions.

$$(x_i - x_j)^2 + (y_i - y_j)^2 + (z_i - z_j)^2 = d_{ij}^2 \text{ for } ij \in E$$

- # complex solutions is an upper bound
- Loose mixed volume bound

Sphere Equations

$$x_i^2 + y_i^2 + z_i^2 = s_i \text{ for } i \in V$$

$$s_i + s_j - 2(x_i x_j + y_i y_j + z_i z_j) = d_{ij}^2 \text{ for } ij \in E$$

Cayley-Menger matrix

$$CM = \begin{pmatrix} 0 & 1 & 1 & \cdots & 1 \\ 1 & 0 & d_{12}^2 & \cdots & d_{1n}^2 \\ 1 & d_{12}^2 & 0 & \ddots & \cdots \\ \cdots & \cdots & \ddots & \ddots & \cdots \\ 1 & d_{1n}^2 & d_{2n}^2 & \cdots & 0 \end{pmatrix}$$

Theorem (Cayley, Menger)

The distances of a CM matrix are embeddable in \mathbb{R}^D iff

- *$\text{rank}(CM) = D + 2$, and*
- *$(-1)^k \det(CM') \geq 0$, for every submatrix CM' with size $k + 1 \leq D + 2$ that includes the first row and column.*

Distance Geometry

Cayley-Menger matrix

$$CM = \begin{pmatrix} 0 & 1 & 1 & \cdots & 1 \\ 1 & 0 & d_{12}^2 & \cdots & d_{1n}^2 \\ 1 & d_{12}^2 & 0 & \ddots & \cdots \\ \cdots & \cdots & \ddots & \ddots & \cdots \\ 1 & d_{1n}^2 & d_{2n}^2 & \cdots & 0 \end{pmatrix}$$


Theorem (Cayley, Menger)

The distances of a CM matrix are embeddable in \mathbb{R}^3 iff

- *$\text{rank}(CM) = 5$, and*
- *positivity, triangular and tetrahedral inequalities must be satisfied.*

Distance geometry subsystems – Example

$$\begin{pmatrix} 0 & 1 & 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 0 & d_{12}^2 & d_{13}^2 & d_{14}^2 & d_{15}^2 & d_{16}^2 & x_1 \\ 1 & d_{21}^2 & 0 & d_{23}^2 & x_2 & x_3 & d_{26}^2 & d_{27}^2 \\ 1 & d_{31}^2 & d_{32}^2 & 0 & d_{34}^2 & x_4 & x_5 & d_{37}^2 \\ 1 & d_{41}^2 & x_2 & d_{43}^2 & 0 & d_{45}^2 & x_6 & d_{47}^2 \\ 1 & d_{51}^2 & x_3 & x_4 & d_{54}^2 & 0 & d_{56}^2 & d_{57}^2 \\ 1 & d_{61}^2 & d_{62}^2 & x_5 & x_6 & d_{65}^2 & 0 & d_{67}^2 \\ 1 & x_1 & d_{72}^2 & d_{73}^2 & d_{74}^2 & d_{75}^2 & d_{76}^2 & 0 \end{pmatrix}$$



- 21 equations in 6 variables
- Every solution of 3x3 subsystem corresponds to a unique embedding
- Eliminate two more variables using resultants

Homotopy Continuation

- PHCpack (Verschelde, 2014)
- Starting system based on structure of equations

RootFinding package (Maple 18)


- Isolate (Rouillier, 1999, Rouillier, Zimmermann, 2003, Aubry, Lazard, Moreno Maza, 1999, Xia, Yang, 2002)
- Algebraic sets over \mathbb{R}

Construction of Geiringer graphs



- H1 and H2 steps always rigid (sufficient for $|V| \leq 12$)
- H1 steps double the number of embeddings


Construction of Geiringer graphs



- H1 and H2 steps always rigid (sufficient for $|V| \leq 12$)
- H1 steps double the number of embeddings
- Known number of real embeddings for $|V| \leq 6$ (Emiris, Mourrain, 1999)


Geiringer graphs with 7 vertices

The graphs that cannot be constructed by H1 in the last step:



	G_{48}	G_{32a}	G_{32b}	G_{24}	G_{16a}	G_{16b}
MV of sphere eq.	48	32	32	32	32	32
MV of dist. subsyst.	48	32	32	24	24	16
# complex emb.	48	32	32	24	16	16

Geiringer graphs with 8 vertices



	G_{128}	G_{160}
MV of sphere eq.	128	160
MV of dist. subsyst.	128	160
# complex emb.	128	160

Maximizing the number of real embeddings

Goal

Find edge lengths with as many real solutions as possible

Maximizing the number of real embeddings

Goal

Find edge lengths with as many real solutions as possible

- Local methods
 - ▶ Stochastic (7-vertex Laman graph – Emiris, Moroz, 2011)
 - ▶ Gradient descent (Stewart-Gough platform – Dietmaier, 1998)
- Global methods
 - ▶ Huge size of parameter space
 - ▶ `RootFinding[Parametric]`
(Liang, Gerhard, Jeffrey, Moroz, 2009)

Maximizing the number of real embeddings

Goal


Find edge lengths with as many real solutions as possible

- Local methods
 - ▶ Stochastic (7-vertex Laman graph – Emiris, Moroz, 2011)
 - ▶ Gradient descent (Stewart-Gough platform – Dietmaier, 1998)
- Global methods
 - ▶ Huge size of parameter space
 - ▶ `RootFinding[Parametric]`
(Liang, Gerhard, Jeffrey, Moroz, 2009)
- Global search over subset of parameters
 - ▶ Coupler curves (6-vertex Laman graph – Borcea, Streinu, 2004)

Coupler Curves

Removing an edge uc from a Geiringer graph breaks rigidity.


The curve traced by the vertex c is called a *coupler curve*.



Coupler Curves


Removing an edge uc from a Geiringer graph breaks rigidity.

The curve traced by the vertex c is called a *coupler curve*.




The real embeddings of G correspond to the intersections of the coupler curve with the sphere centered at u with radius d_{uc} .


Invariance of coupler curve



Invariance of coupler curve




Invariance of coupler curve




- The coupler curve of c is invariant to the position of u
- 2-parameter family changing 4 edge lengths
- Increasing the number of real embeddings


Example



Example




Sampling




Sampling

If wc is an edge, then the coupler curve of c is a spherical curve.




Sampling

If wc is an edge, then the coupler curve of c is a spherical curve.



- Find φ and θ that maximize the number of real solutions
- Repeat the procedure with another suitable subgraph

Results



	G_{48}	G_{32a}	G_{32b}	G_{24}	G_{16a}	G_{16b}	G_{128}	G_{160}
# compl.	48	32	32	24	16	16	128	160
# real	48	32	32	24	16	16	128	≥ 132

Source code & results: jan.legersky.cz/project/spatialgraphembeddings/

Asymptotic bounds

Let n be the number of vertices.

- The number of real embeddings is at most

$$\frac{2^{n-3}}{n-2} \binom{3n-6}{n-3} \quad (\text{Borcea, Streinu, 2004})$$


which behaves asymptotically as 8^n

- There are graphs with $\lfloor 2.51984^n \rfloor$ real embeddings (Emiris, Tsigaridas, Varvitsiotis, 2013)
- There are graphs with $\lfloor 3.0682^n \rfloor$ complex embeddings (Grasegger, Koutschan, Tsigaridas, 2018)

Lower bound on the maximum number of real embeddings

Theorem

There are graphs with $\lfloor 2.6553^n \rfloor$ real embeddings.



There are at least 132^k real embeddings, where k is the number of copies of G_{160} . For a graph with n vertices, $k = \lfloor \frac{n-3}{5} \rfloor$.

Future work

- Tight bound for the number of real embeddings of G_{160}
- Number of real embeddings of 9-vertex Geiringer graphs
- Improving lower bounds
- Other dimensions

Thank you

vbartzos@di.uoa.gr
users.uoa.gr/~vbartzos

emiris@di.uoa.gr
cgi.di.uoa.gr/~emiris

jan.legersky@risc.jku.at
jan.legersky.cz

elias.tsigaridas@inria.fr
polsys.lip6.fr/~elias