

Computing Hypergeometric Solutions of Second Order Linear Differential Equations using Quotients of Formal Solutions

Erdal Imamoglu & Mark van Hoeij

Florida State University
Department of Mathematics

University of Bath
ISSAC 2015
July 8, 2015

Outline of the Talk

1. Introduction and Riemann-Hurwitz Type Formula
2. Algorithm Outline and Examples

1. Introduction and Riemann-Hurwitz Type Formula

${}_2F_1$ -type Solutions

Let a second order differential equation $L_{inp}(y) = 0$ be given.

Here $L_{inp} \in \mathbb{C}(x)[\partial]$, $\text{ord}(L_{inp}) = 2$, $\partial = \frac{d}{dx}$.

${}_2F_1$ -type Solutions

Let a second order differential equation $L_{inp}(y) = 0$ be given.

Here $L_{inp} \in \mathbb{C}(x)[\partial]$, $\text{ord}(L_{inp}) = 2$, $\partial = \frac{d}{dx}$.

We will give an algorithm to find ${}_2F_1$ -type solutions of L_{inp} in the form of

$$\exp\left(\int r dx\right) {}_2F_1(a_1, a_2; b_1; f)$$

where $a_1, a_2, b_1 \in \mathbb{Q}$ and $r, f \in \overline{\mathbb{Q}(x)}$.

${}_2F_1$ -type Solutions

Let a second order differential equation $L_{inp}(y) = 0$ be given.

Here $L_{inp} \in \mathbb{C}(x)[\partial]$, $\text{ord}(L_{inp}) = 2$, $\partial = \frac{d}{dx}$.

We will give an algorithm to find ${}_2F_1$ -type solutions of L_{inp} in the form of

$$\exp\left(\int r dx\right) {}_2F_1(a_1, a_2; b_1; f)$$

where $a_1, a_2, b_1 \in \mathbb{Q}$ and $r, f \in \overline{\mathbb{Q}(x)}$.

We need to find

- 1 $a_1, a_2, b_1 \in \mathbb{Q}$,
- 2 Pullback function $f \in \overline{\mathbb{Q}(x)}$,
- 3 $r \in \overline{\mathbb{Q}(x)}$.

Base Operator L_B

If solutions of a second order differential operator L at $x = 0$ are,

$$y_1 = x^0 + \dots,$$

$$y_2 = x^{\frac{1}{3}} + \dots,$$

then we say that *exponent difference* of L at $x = 0$ is

$$\Delta(L, 0) := \left| 0 - \frac{1}{3} \right|.$$

Base Operator L_B

If solutions of a second order differential operator L at $x = 0$ are,

$$y_1 = x^0 + \dots,$$

$$y_2 = x^{\frac{1}{3}} + \dots,$$

then we say that *exponent difference* of L at $x = 0$ is

$$\Delta(L, 0) := |0 - \frac{1}{3}|.$$

Hypergeometric function ${}_2F_1(a_1, a_2; b_1; x)$ is a solution of a Gauss Hypergeometric Differential Operator (GHDO)

$$L_B = x(1-x)\partial^2 + (b_1 - (a_1 + a_2 + 1)x)\partial - a_1a_2$$

at $x = 0$.

Base Operator L_B

If solutions of a second order differential operator L at $x = 0$ are,

$$y_1 = x^0 + \dots,$$

$$y_2 = x^{\frac{1}{3}} + \dots,$$

then we say that *exponent difference* of L at $x = 0$ is

$$\Delta(L, 0) := |0 - \frac{1}{3}|.$$

Hypergeometric function ${}_2F_1(a_1, a_2; b_1; x)$ is a solution of a Gauss Hypergeometric Differential Operator (GHDO)

$$L_B = x(1-x)\partial^2 + (b_1 - (a_1 + a_2 + 1)x)\partial - a_1a_2$$

at $x = 0$.

L_B has three singularities $\{0, 1, \infty\}$ with exponent differences

$$\alpha_0 = |1 - b_1|, \quad \alpha_1 = |b_1 - a_1 - a_2|, \quad \alpha_\infty = |a_1 - a_2|.$$

Base Operator L_B

If solutions of a second order differential operator L at $x = 0$ are,

$$y_1 = x^0 + \dots,$$

$$y_2 = x^{\frac{1}{3}} + \dots,$$

then we say that *exponent difference* of L at $x = 0$ is

$$\Delta(L, 0) := |0 - \frac{1}{3}|.$$

Hypergeometric function ${}_2F_1(a_1, a_2; b_1; x)$ is a solution of a Gauss Hypergeometric Differential Operator (GHDO)

$$L_B = x(1-x)\partial^2 + (b_1 - (a_1 + a_2 + 1)x)\partial - a_1a_2$$

at $x = 0$.

L_B has three singularities $\{0, 1, \infty\}$ with exponent differences

$$\alpha_0 = |1 - b_1|, \quad \alpha_1 = |b_1 - a_1 - a_2|, \quad \alpha_\infty = |a_1 - a_2|.$$

If we can find $\alpha_0, \alpha_1, \alpha_\infty$, then we will have a_1, a_2, b_1 .

Set of Candidates for $\alpha_0, \alpha_1, \alpha_\infty$

Assume L_{inp} has a solution in the form

$$\exp\left(\int r dx\right) {}_2F_1(a_1, a_2; b_1; f).$$

Set of Candidates for $\alpha_0, \alpha_1, \alpha_\infty$

Assume L_{inp} has a solution in the form

$$\exp\left(\int r dx\right) {}_2F_1(a_1, a_2; b_1; f).$$

This implies:

$$L_B \xrightarrow{f} C \quad M \xrightarrow{r} E \quad L_{inp}$$
$$y_i(x) \mapsto y_i(f) \mapsto \exp\left(\int r dx\right) y_i(f), \quad i = 1, 2.$$

Set of Candidates for $\alpha_0, \alpha_1, \alpha_\infty$

Assume L_{inp} has a solution in the form

$$\exp\left(\int r dx\right) {}_2F_1(a_1, a_2; b_1; f).$$

This implies:

$$L_B \xrightarrow{f} C \quad M \xrightarrow{r} E \quad L_{inp}$$
$$y_i(x) \mapsto y_i(f) \mapsto \exp\left(\int r dx\right) y_i(f), \quad i = 1, 2.$$

	L_B	L_{inp}
singularities	$0, 1, \infty$	p_1, \dots, p_r
exponent differences	$\alpha_0, \alpha_1, \alpha_\infty$	$\Delta_{p_1}, \dots, \Delta_{p_r}$

Set of Candidates for $\alpha_0, \alpha_1, \alpha_\infty$

Assume L_{inp} has a solution in the form

$$\exp\left(\int r dx\right) {}_2F_1(a_1, a_2; b_1; f).$$

This implies:

$$L_B \xrightarrow{f} C \quad M \xrightarrow{r} E \quad L_{inp}$$
$$y_i(x) \mapsto y_i(f) \mapsto \exp\left(\int r dx\right) y_i(f), \quad i = 1, 2.$$

	L_B	L_{inp}
singularities	$0, 1, \infty$	p_1, \dots, p_r
exponent differences	$\alpha_0, \alpha_1, \alpha_\infty$	$\Delta_{p_1}, \dots, \Delta_{p_r}$

$$\alpha_0, \alpha_1, \alpha_\infty \in \left\{ \frac{a}{b} : a \in \{1, \Delta_{p_1}, \dots, \Delta_{p_r}\}, 1 \leq b \leq \deg(f) \right\}$$

Set of Candidates for $\alpha_0, \alpha_1, \alpha_\infty$

Assume L_{inp} has a solution in the form

$$\exp\left(\int r dx\right) {}_2F_1(a_1, a_2; b_1; f).$$

This implies:

$$L_B \xrightarrow{f} C \quad M \xrightarrow{r} E \quad L_{inp}$$
$$y_i(x) \mapsto y_i(f) \mapsto \exp\left(\int r dx\right) y_i(f), \quad i = 1, 2.$$

	L_B	L_{inp}
singularities	$0, 1, \infty$	p_1, \dots, p_r
exponent differences	$\alpha_0, \alpha_1, \alpha_\infty$	$\Delta_{p_1}, \dots, \Delta_{p_r}$

$$\alpha_0, \alpha_1, \alpha_\infty \in \left\{ \frac{a}{b} : a \in \{1, \Delta_{p_1}, \dots, \Delta_{p_r}\}, 1 \leq b \leq \deg(f) \right\}$$

Problem: This set is finite but **too large**.

Differential Operators on Algebraic Curves

L_B and L_{inp} are in $\mathbb{C}(x)[\partial]$, i.e., they are defined on \mathbb{P}^1 .

Differential Operators on Algebraic Curves

L_B and L_{inp} are in $\mathbb{C}(x)[\partial]$, i.e., they are defined on \mathbb{P}^1 .

In general:

Let X be an algebraic curve.

Let $\mathbb{C}(X)$ be its function field.

$D_{\mathbb{C}(X)} := \mathbb{C}(X)[\partial_t]$ where $t \in \mathbb{C}(X)$ with $t' \neq 0$.

$L \in D_{\mathbb{C}(X)}$ is a differential operator.

Riemann-Hurwitz Type Formula for Differential Equations

Theorem

Let X, Y be algebraic curves and $\mathbb{C}(X), \mathbb{C}(Y)$ be their function fields.

Riemann-Hurwitz Type Formula for Differential Equations

Theorem

Let X, Y be algebraic curves and $\mathbb{C}(X), \mathbb{C}(Y)$ be their function fields.

Let $f : X \rightarrow Y$ be a non-constant morphism.

Riemann-Hurwitz Type Formula for Differential Equations

Theorem

Let X, Y be algebraic curves and $\mathbb{C}(X), \mathbb{C}(Y)$ be their function fields.

Let $f : X \rightarrow Y$ be a non-constant morphism.

f corresponds a homomorphism $\mathbb{C}(Y) \rightarrow \mathbb{C}(X)$.

This induces a homomorphism $D_{\mathbb{C}(Y)} \rightarrow D_{\mathbb{C}(X)}$.

Riemann-Hurwitz Type Formula for Differential Equations

Theorem

Let X, Y be algebraic curves and $\mathbb{C}(X), \mathbb{C}(Y)$ be their function fields.

Let $f : X \rightarrow Y$ be a non-constant morphism.

f corresponds a homomorphism $\mathbb{C}(Y) \rightarrow \mathbb{C}(X)$.

This induces a homomorphism $D_{\mathbb{C}(Y)} \rightarrow D_{\mathbb{C}(X)}$.

If $L_1 \in D_{\mathbb{C}(Y)}$ with $\text{ord}(L_1)=2$ and

L_2 is the corresponding element in $D_{\mathbb{C}(X)}$, then

$$2 - 2g_X + \Delta^{\Sigma^{-1}}(L_2) = \deg(f) \left(2 - 2g_Y + \Delta^{\Sigma^{-1}}(L_1) \right)$$

where g_X, g_Y are genus of X, Y resp., and

$$\Delta^{\Sigma^{-1}}(L) := \sum_p (\Delta(L, p) - 1),$$

$\Delta(L, p) :=$ exponent difference of L at $x = p$.

Riemann-Hurwitz Type Formula for Our Case

In our case $X = Y = \mathbb{P}^1$, so $g_X = g_Y = 0$.

Riemann-Hurwitz Type Formula for Our Case

In our case $X = Y = \mathbb{P}^1$, so $g_X = g_Y = 0$.

$$L_B \xrightarrow{f} C \quad M \xrightarrow{r} E \quad L_{\text{inp}}$$
$$y_i(x) \mapsto y_i(f) \mapsto \exp\left(\int r dx\right) y_i(f), \quad i = 1, 2.$$

where L_B has exponent differences $[\alpha_0, \alpha_1, \alpha_\infty]$ at $\{0, 1, \infty\}$.

Riemann-Hurwitz Type Formula for Our Case

In our case $X = Y = \mathbb{P}^1$, so $g_X = g_Y = 0$.

$$L_B \xrightarrow{f} L_C \quad M \xrightarrow{r} L_{\text{inp}}$$
$$y_i(x) \mapsto y_i(f) \mapsto \exp\left(\int r dx\right) y_i(f), \quad i = 1, 2.$$

where L_B has exponent differences $[\alpha_0, \alpha_1, \alpha_\infty]$ at $\{0, 1, \infty\}$.

\xrightarrow{r} **does not affect exponent differences.**

Riemann-Hurwitz Type Formula for Our Case

In our case $X = Y = \mathbb{P}^1$, so $g_X = g_Y = 0$.

$$L_B \xrightarrow{f} M \xrightarrow{r} L_{inp}$$
$$y_i(x) \mapsto y_i(f) \mapsto \exp\left(\int r dx\right) y_i(f), \quad i = 1, 2.$$

where L_B has exponent differences $[\alpha_0, \alpha_1, \alpha_\infty]$ at $\{0, 1, \infty\}$.

\xrightarrow{r} **does not affect exponent differences.**

Therefore:

$$2 + \Delta^{\Sigma^{-1}}(L_{inp}) = \deg(f) \left(2 + \Delta^{\Sigma^{-1}}(L_B) \right)$$

Riemann-Hurwitz Type Formula for Our Case

In our case $X = Y = \mathbb{P}^1$, so $g_X = g_Y = 0$.

$$L_B \xrightarrow{f} M \xrightarrow{r} L_{in p}$$
$$y_i(x) \mapsto y_i(f) \mapsto \exp\left(\int r dx\right) y_i(f), \quad i = 1, 2.$$

where L_B has exponent differences $[\alpha_0, \alpha_1, \alpha_\infty]$ at $\{0, 1, \infty\}$.

\xrightarrow{r} **does not affect exponent differences.**

Therefore:

$$2 + \Delta^{\Sigma^{-1}}(L_{in p}) = \deg(f) \left(2 + \Delta^{\Sigma^{-1}}(L_B) \right)$$

Explicitly:

$$2 + \sum_{p \in \mathbb{P}^1} (\Delta(L_{in p}, p) - 1) = \deg(f) \left(2 + \sum_{i \in \{0, 1, \infty\}} (\alpha_i - 1) \right)$$

Computing Candidates for $\alpha_0, \alpha_1, \alpha_\infty$

	L_B	L_{inp}
singularities	$0, 1, \infty$	p_1, \dots, p_r
exponent differences	$\alpha_0, \alpha_1, \alpha_\infty$	$\Delta_{p_1}, \dots, \Delta_{p_r}$

$$\alpha_0, \alpha_1, \alpha_\infty \in \left\{ \frac{a}{b} : a \in \{1, \Delta_{p_1}, \dots, \Delta_{p_r}\}, 1 \leq b \leq \deg(f) \right\}$$

This set is finite but too large.

Computing Candidates for $\alpha_0, \alpha_1, \alpha_\infty$

	L_B	L_{inp}
singularities	$0, 1, \infty$	p_1, \dots, p_r
exponent differences	$\alpha_0, \alpha_1, \alpha_\infty$	$\Delta_{p_1}, \dots, \Delta_{p_r}$

$$\alpha_0, \alpha_1, \alpha_\infty \in \left\{ \frac{a}{b} : a \in \{1, \Delta_{p_1}, \dots, \Delta_{p_r}\}, 1 \leq b \leq \deg(f) \right\}$$

This set is finite but too large.

Formula

$$2 + \sum_{p \in \mathbb{P}^1} (\Delta(L_{inp}, p) - 1) = \deg(f) \left(2 + \sum_{i \in \{0, 1, \infty\}} (\alpha_i - 1) \right)$$

eliminates **vast majority** of the candidates.

2. Algorithm Outline and Examples

Algorithm Outline

INPUT:

A second order differential operator $L_{inp} \in \mathbb{C}(x)[\partial]$.

Algorithm Outline

INPUT:

A second order differential operator $L_{inp} \in \mathbb{C}(x)[\partial]$.

OUTPUT:

A basis $[\text{Sol}_1, \text{Sol}_2]$ of ${}_2F_1$ -type solutions of L_{inp} in the form of

$$\exp\left(\int r dx\right) {}_2F_1(a_1, a_2; b_1; f)$$

or

an empty list [].

Algorithm Outline

INPUT:

A second order differential operator $L_{inp} \in \mathbb{C}(x)[\partial]$.

OUTPUT:

A basis $[\text{Sol}_1, \text{Sol}_2]$ of ${}_2F_1$ -type solutions of L_{inp} in the form of

$$\exp\left(\int r dx\right) {}_2F_1(a_1, a_2; b_1; f)$$

or

an empty list $[\]$.

Note: If $f \in \overline{\mathbb{Q}(x)} \setminus \mathbb{Q}(x)$, then additional input is required.

STEP 1: Candidate Exponent Differences

Compute the list of candidate exponent differences

$$[\alpha_0, \alpha_1, \alpha_\infty, d]$$

for candidate base operators L_B .

If there is no candidate, then return [].

Algorithm Outline

STEP 2: Quotient Method

Choose a candidate $[\alpha_0, \alpha_1, \alpha_\infty, d]$, compute the corresponding operator L_B .

Algorithm Outline

STEP 2: Quotient Method

Choose a candidate $[\alpha_0, \alpha_1, \alpha_\infty, d]$, compute the corresponding operator L_B .

If L_{inp} has a ${}_2F_1$ -type solution coming from L_B , then

$$L_B \xrightarrow{f} C \xrightarrow{r} E L_{inp} \quad (1)$$

Algorithm Outline

STEP 2: Quotient Method

Choose a candidate $[\alpha_0, \alpha_1, \alpha_\infty, d]$, compute the corresponding operator L_B .

If L_{inp} has a ${}_2F_1$ -type solution coming from L_B , then

$$L_B \xrightarrow{f} C \xrightarrow{r} E L_{inp} \quad (1)$$

Compute formal solutions of y_1, y_2 of L_B at s , and Y_1, Y_2 of L_{inp} at $f(s)$ up to order a .

Algorithm Outline

STEP 2: Quotient Method

Choose a candidate $[\alpha_0, \alpha_1, \alpha_\infty, d]$, compute the corresponding operator L_B .

If L_{inp} has a ${}_2F_1$ -type solution coming from L_B , then

$$L_B \xrightarrow{f} C \xrightarrow{r} E L_{inp} \quad (1)$$

Compute formal solutions of

y_1, y_2 of L_B at s , and Y_1, Y_2 of L_{inp} at $f(s)$ up to order a .

If (1) is true, then

$$y_i(x) \mapsto Y_i(x) = \exp\left(\int r dx\right)y_i(f), \quad i = 1, 2.$$

Algorithm Outline

STEP 2: Quotient Method

Choose a candidate $[\alpha_0, \alpha_1, \alpha_\infty, d]$, compute the corresponding operator L_B .

If L_{inp} has a ${}_2F_1$ -type solution coming from L_B , then

$$L_B \xrightarrow{f} c \xrightarrow{r} E L_{inp} \quad (1)$$

Compute formal solutions of

y_1, y_2 of L_B at s , and Y_1, Y_2 of L_{inp} at $f(s)$ up to order a .

If (1) is true, then

$$y_i(x) \mapsto Y_i(x) = \exp\left(\int r dx\right) y_i(f), \quad i = 1, 2.$$

Let $q = \frac{y_1}{y_2}$ and $Q = \frac{Y_1}{Y_2}$. Then

$$q(f) = c \cdot Q(x) \implies f = q^{-1}(c \cdot Q(x))$$

for an unknown constant c .

Algorithm Outline

STEP 2 (continued):

There are infinitely many choices for c .

Algorithm Outline

STEP 2 (continued):

There are infinitely many choices for c .

Idea: Choose a suitable prime p and compute

$$f \equiv q^{-1}(c \cdot Q(f)) \pmod{(x^{a-1}, p)}$$

by looping $c = 1, \dots, p - 1$.

For each c try Rational Function Reconstruction to find $f \in \mathbb{F}_p[x]$.

Algorithm Outline

STEP 2 (continued):

There are infinitely many choices for c .

Idea: Choose a suitable prime p and compute

$$f \equiv q^{-1}(c \cdot Q(f)) \pmod{(x^{a-1}, p)}$$

by looping $c = 1, \dots, p - 1$.

For each c try Rational Function Reconstruction to find $f \in \mathbb{F}_p[x]$.

If this succeeds, then try p -adic Hensel lifting
and Rational Number Reconstruction to find $f \in \mathbb{Q}(x)$.

STEP 2 (continued):

For algebraic pullback functions:

Compute the minimal polynomial of $f \bmod (x^{a-1}, p)$.

Hensel lift it, and try Rational Number Reconstruction.

STEP 3:

Compute the operator M such that

$$L_B \xrightarrow{f} C \quad M \xrightarrow{r} E \quad L_{inp}$$

Then use existing algorithms to find r .

STEP 3:

Compute the operator M such that

$$L_B \xrightarrow{f} C \quad M \xrightarrow{r} E \quad L_{inp}$$

Then use existing algorithms to find r .

Compute solutions of L_B and apply transformations to them.

STEP 3:

Compute the operator M such that

$$L_B \xrightarrow{f} C \quad M \xrightarrow{r} E \quad L_{inp}$$

Then use existing algorithms to find r .

Compute solutions of L_B and apply transformations to them.

Return a basis $[\text{Sol}_1, \text{Sol}_2]$ of ${}_2F_1$ -type solutions of L_{inp} .

Examples

```
> L := Linp[99];
      L := 400 (7 x^2 - 7 x + 8) (x + 7) (x + 8) Dx^2 + (7560 x^3 + 85120 x^2 + 197560 x - 98880) Dx + 1323 x^2 + 11270 x + 12803
```

```
> s := find_2fl(L);
```

```
[x, x, x]
[1/5, 1/5, 1/4, 2]
13, 1
[2/15, 2/15, 1/2, 3]
23, 1
[1/15, 1/5, 1/2, 3]
13, 1
[1/5, 1/2, 1/15, 3]
13, 1
[1/5, 1/2, 1/8, 4]
13, 1
[1/5, 1/5, 1/2, 7]
13, 1
13, 2
13, 3
13, 4
13, 5
13, 6
13, 7
13, 8
13, 9
```

$$s := \frac{\text{hypergeom}\left(\left[\frac{1}{20}, \frac{11}{20}\right], \left[\frac{4}{5}\right], \frac{(x+7)^2 x^5}{(x+8)((x+7)^3 - 18(x+7)^2 + 101x + 547)^2}\right)}{\left((x+7)^3 - 18(x+7)^2 + 101x + 547\right)^{1/10} (x+8)^{1/20}} \cdot \frac{\left(\frac{(x+7)^2 x^5}{(x+8)((x+7)^3 - 18(x+7)^2 + 101x + 547)^2}\right)^{1/5} \text{hypergeom}\left(\left[\frac{1}{4}, \frac{3}{4}\right], \left[\frac{6}{5}\right], \frac{(x+7)^2 x^5}{(x+8)((x+7)^3 - 18(x+7)^2 + 101x + 547)^2}\right)}{\left((x+7)^3 - 18(x+7)^2 + 101x + 547\right)^{1/10} (x+8)^{1/20}}$$

```
> seq( simplify( eval( diffop2de(L,y(x)), y(x)=i ) ), i=s );
      0, 0
```

```
> find_expdiffs( singDiffs(L) );
```

```
[x, x, x]
```

```
[[1/5, 1/5, 1/4, 2], [2/15, 2/15, 1/2, 3], [1/15, 1/5, 1/2, 3], [1/8, 1/5, 1/2, 4], [1/5, 1/5, 1/2, 7], [1/5, 1/4, 1/2, 14]]
```

```

> L := LLinp[92];
L := Dx^2 - 1/16 * (65028096 x^8 - 140894208 x^7 + 18047232 x^6 + 92111232 x^5 - 14986448 x^4 - 15758239 x^3 - 3896340 x^2 - 744800 x - 78400) / ((x-1)^2 (21x+4)^2 (144x^2-84x-35)^2)

```

$$L := Dx^2 - \frac{1}{16} \frac{65028096 x^8 - 140894208 x^7 + 18047232 x^6 + 92111232 x^5 - 14986448 x^4 - 15758239 x^3 - 3896340 x^2 - 744800 x - 78400}{(x-1)^2 (21x+4)^2 (144x^2-84x-35)^2}$$

```

> s := find_2f1(L);
[0, 0, x]
[0, x, x]
[0, 1/3, 1/2, 7]
23, 1
23, 2
23, 3
23, 4
23, 5
23, 6
23, 7
s := \frac{(x-1)^{1/6} \sqrt{x} \sqrt{144x^2-84x-35} (21x+4)^{1/6} \operatorname{hypergeom}\left(\left[\frac{1}{12}, \frac{7}{12}\right], \left[\frac{2}{3}\right], 1 - \frac{x^3(144x^2-84x-35)^2}{(x-1)(144x^3-12x^2-23x-4)^2}\right)}{(144x^3-12x^2-23x-4)^{1/6}},
\frac{1}{(144x^3-12x^2-23x-4)^{1/6}} \left( (x-1)^{1/6} \sqrt{x} \sqrt{144x^2-84x-35} (21x+4)^{1/6} \operatorname{hypergeom}\left(\left[\frac{5}{12}, \frac{11}{12}\right], \left[\frac{4}{3}\right], 1 - \frac{x^3(144x^2-84x-35)^2}{(x-1)(144x^3-12x^2-23x-4)^2}\right) \right) \left( \frac{x^3(144x^2-84x-35)^2}{(x-1)(144x^3-12x^2-23x-4)^2} - 1 \right)^{1/3} \right)

```

$$s := \frac{(x-1)^{1/6} \sqrt{x} \sqrt{144x^2-84x-35} (21x+4)^{1/6} \operatorname{hypergeom}\left(\left[\frac{1}{12}, \frac{7}{12}\right], \left[\frac{2}{3}\right], 1 - \frac{x^3(144x^2-84x-35)^2}{(x-1)(144x^3-12x^2-23x-4)^2}\right)}{(144x^3-12x^2-23x-4)^{1/6}},$$

$$\frac{1}{(144x^3-12x^2-23x-4)^{1/6}} \left((x-1)^{1/6} \sqrt{x} \sqrt{144x^2-84x-35} (21x+4)^{1/6} \operatorname{hypergeom}\left(\left[\frac{5}{12}, \frac{11}{12}\right], \left[\frac{4}{3}\right], 1 - \frac{x^3(144x^2-84x-35)^2}{(x-1)(144x^3-12x^2-23x-4)^2}\right) \right) \left(\frac{x^3(144x^2-84x-35)^2}{(x-1)(144x^3-12x^2-23x-4)^2} - 1 \right)^{1/3} \right)$$

```

> seq( simplify( eval( diffop2de(L,y(x)), y(x)=i) ), i=s );
0, 0
> find_expdiffs( singDiffs(L) );
[0, 0, x]
[0, x, x]

```

$$\left[\left[0, \frac{1}{3}, \frac{1}{2}, 7 \right] \right]$$

Thank You